ACCELERATING OPTICAL BEAMS

Accelerating Beyond the Horizon

R esearch on accelerating beams has developed rapidly since 2007.¹ An ideal paraxial accelerating beam follows a parabolic trajectory while preserving its amplitude structure indefinitely as waves emitted from all points interfere to a propagationinvariant structure shifting laterally on a curved trajectory. This phenomenon has led to many ideas, including particle guidance along curves and nonlinear shape-preserving accelerating beams.

Until this year, however, the only shapepreserving accelerating solutions ever found were derived from the paraxial wave equation, which gave an Airy beam propagating on a parabolic trajectory. This solution is limited to small angles: After some distance, an Airy beam trajectory will inevitably reach a steep angle where the dynamics are no longer shape-preserving. Similarly, the paraxial regime cannot describe accelerating beams with narrow lobes—comparable to the optical wavelength—whose steep bending occurs within tens of wavelengths. Previous attempts to find non-paraxial accelerating beams showed deformation and breakup.

In a recent paper,¹ this apparent limit was tackled through first-principles analysis, starting from Maxwell's equations.² This theory found a fundamentally new solution to the Maxwell equations corresponding to nondiffracting spatially accelerating beams along a circular trajectory, exhibiting shapepreserving bending with the Poynting vector of the main lobe displaying a turn of close to 180° (with an initial tilt).² Theory was followed by experimental confirmation.³ The figure depicts experimental and theoretical results. The online video shows formation of a shape-preserving beam propagating at a circular trajectory, bending by almost 90°; the beam results from interference among sequentially added lobes with appropriate amplitude and phase.



(a) Experiments (left) and theory (right) depicting the intensity of a shape-preserving beam bending at more than 90° degrees. (b) The use of accelerating beams also allows novel applications such as femtosecond-pulsed beams propagating on circular trajectories to fabricate micron-scale curved structures in photonic materials such as silicon.

Interestingly, any circular trajectory can support a family of accelerating solutions, whereby their superpositions form periodic accelerating "breathers." In scalar form, these beams are exact solutions for non-dispersive accelerating wavepackets of the common (Helmholtz-type) wave equation describing time-harmonic waves. Hence, this work has implications for many waves in nature, ranging from acoustic and elastic to surface waves in fluids and membranes. It shows that exact non-diffracting beams are no longer necessarily only straight-line Bessel-type beams as believed since Stratton's classic 1941 text on electromagnetism.⁴ The family has now been extended to include self-bending beams.²

This work generated much follow up, with extensions to nonlinear media, non-paraxial and non-circular trajectories, 3-D accelerating beams with trajectories that do not lie in a single plane, and even technical applications showing material processing with curved features. It also brings the physics of accelerating beams into the regime of super-resolution, through the sub-wavelength features of the solutions. **OPN**



Visit **www.opnmagazine-digital.com** to view the video that accompanies this article.

Researchers

Ido Kaminer, Elad Greenfield, Rivka Bekenstein, Jonathan Nemirovsky and Mordechai Segev (msegev@techunix. technion.ac.il) Department of physics and Solid State Institute, Technion, Haifa, Israel Amaury Mathis,

Luc Froehly, François Courvoisier and John M. Dudley Département d'Optique Institut FEMT0-ST, Université de Franche-Comté, Besançon, France

References

- 1. G.A. Siviloglou and D.N. Christodoulides. Phys. Rev. Lett. **99**, 213901 (2007).
- I. Kaminer et al. Phys. Rev. Lett. 108, 163901 (2012).
- F. Courvoisier et al. Opt. Lett. 37, 1736-1738 (2012).
- J.A. Stratton. Electromagnetic Theory, 1941.